

FILE 'CAPLUS' ENTERED AT 08:36:16 ON 11 DEC 2002
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGE/TERMS" FOR DETAILS.
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> s antibody?

L1 2374153 ANTIBOD.

=> s 11 and "GPVI"

MISMATCHED QUOTE 'AND "GPVI"

Quotation marks or apostrophes must be used in pairs,
one before and one after the expression you are setting
off or masking.

=> s 11 and "GPVI"

L1 167 11 AND "GPVI"

=> s 12 and monoclonal

L1 91 12 AND MONOCLONAL

=> s 13 and glycoprotein VI

L1 61 13 AND GLYCOPROTEIN VI

=> s 14 and platelet

L1 61 14 AND PLATELET

=> dup remove 15

PROCESSING COMPLETED FOR L1

L1 11 DUP REMOVE L1 (41 DUPLICATES REMOVED)

=> d 16 1-21 both aka

L1 ANSWER 1 OF 21 MEDLINE

DUPLICATE 1

2002075976 Document Number: 21659770. PubMed ID: 11723134. The

platelet receptor **GPVI** mediates both adhesion and
signaling responses to collagen in a receptor density-dependent fashion.
Chen Hong; Locke Darren; Liu Ying; Liu Changdong; Kahn Mark L. (Department
of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104,
USA.) JOURNAL OF BIOLOGICAL CHEMISTRY, (2002 Jan 25) 277 (4) 3011-9.
Journal code: 2474-111X. ISSN: 0021-9258. Pub. country: United States.
Language: English.

Ab The **platelet** response to collagen is a primary event in
hemostasis and thrombosis, but the precise roles of the numerous
identified **platelet** collagen receptors remain incompletely
defined. Attention has recently focused on **glycoprotein**
VI (**GPVI**), a receptor that is expressed on
platelets in association with a signaling adapter, the Fc receptor
gamma chain (FcRgamma). Genetic and pharmacologic loss of **GPVI**
function results in loss of collagen signaling in **platelets**, but
studies to date have failed to demonstrate that **GPVI**-FcRgamma
expression is sufficient to confer collagen signaling responses. These
results have led to the hypothesis that collagen responses mediated by
GPVI-FcRgamma may require the collagen-binding integrin
alpha2beta1 as a coreceptor, but this model has not been supported by a
recent study of mouse **platelets** lacking alpha2beta1. In the
present study we have used a novel anti-**GPVI** monoclonal
antibody to measure the level of **GPVI** on human
platelets and to guide the development of **GPVI**
-expressing cell lines to assess the role of **GPVI** in mediating
platelet collagen responses. **GPVI** receptor density on
human **platelets** appears tightly regulated, is independent from
the level of alpha2beta1 expression, and significantly exceeds that on

previously characterized **GPVI**-expressing RFL-2H3 cells. Using newly generated **GPVI**-expressing RFL-1H3 cells with receptor sensitivity equivalent to that on human **platelets**, we demonstrate that **GPVI** expression confers both adhesive and signaling response to collagen in a graded fashion that is proportional to the **GPVI** receptor density. These results resolve some of the conflicting data regarding **GPVI**-collagen interactions and demonstrate that 1) **GPVI**-Fc Rgamma expression is sufficient to confer both adhesion and signaling responses to collagen, and 2) **GPVI**-mediated collagen responses are receptor density-dependent at the receptor levels expressed on human **platelets**.

- L6 ANSWER 2 OF 21 MEDLINE DUPLICATE
 2002617267 Document Number: 21174623. Pubmed ID: 12117414. Differential effects of reduced **glycoprotein VI** levels on activation of murine **platelets** by **glycoprotein VI** ligands. Snell Daniel B; Schulte Valerie; Jarvis Gavin E; Arase Kanao; Sakurai Daigo; Saito Takashi; Watson Steve P; Nieuwandt Bernhard. Department of Pharmacology, University of Oxford, Mansfield Road, Oxford OX1 3QT, U.K. • BIOCHEMICAL JOURNAL, 2002 Nov 10 368 (Pt 1) 293-300. Journal code: 00447160. ISSN: 0264-6021. Pub. country: England; United Kingdom. Language: English.
- AB We have investigated the effects of decreased levels of the complex between **glycoprotein VI** (**GPVI**) and the Fc receptor gamma chain (FcRgamma) on responses to collagen and **GPVI**-specific ligands in murine **platelets**. We show that levels of **GPVI**-FcRgamma of the order of 50% and 20% of wild-type levels caused 2- and 5-fold shifts to the right respectively in the dose-response curve for aggregation in response to collagen, the snake toxin convulxin and the monoclonal antibody JQ1. In addition, there is a delay in the onset of aggregation in response to collagen. In contrast, the stimulation of protein tyrosine phosphorylation by collagen (as measured after 150 s) and adhesion to a collagen-coated surface under static conditions were unaffected in **platelets** with 50% and 20% of wild-type levels of **GPVI**. In contrast, responses to a collagen-related peptide (CRP), made up of repeat glycine-proline-hydroxyproline motifs, were markedly inhibited and abolished in **platelets** expressing 50% and 20% of wild-type levels of **GPVI** respectively. We suggest that the marked effect of a reduction in **GPVI** levels on the CRP-induced activation of **platelets** is due to the multivalent nature of CRP and the fact that **GPVI** is its sole receptor on **platelets**. Thus it appears that the interaction of CRP with **GPVI** is determined by a combination of affinity and avidity. The observation that collagen does not behave like CRP in **platelets** expressing reduced levels of **GPVI**, even in the combined presence of blocking antibodies against integrin alpha2beta1 and SPV, suggests that collagen has a greater affinity than CRP for **GPVI**, and/or that other receptors are involved in its binding to **platelets**. The clinical significance of these results is discussed.

- L6 ANSWER 3 OF 21 CAPLAW COPYRIGHT 2002 ACS
 2001:168145 Document No. 134:217.95 Platelet membrane **glycoprotein VI** (**GPVI**) cDNA and protein sequences, and therapeutic uses thereof. Tandon, Narendra; Sun, Bing; Nakamura, Takashi; Yamamoto, Naomasa • Otsuka Pharmaceutical Co., Ltd., Japan). ICT Int. Appl. WO 2001016321 A1 20010307, 71 pp. DESIGNATED STATES: WI: AE, AG, AL, AM, AT, AU, AZ, BA, BE, BG, BR, BY, BS, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, EG, FI, GB, GD, GE, GH, GM, HP, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LF, LR, LS, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AI, AS, BY, KG, KE, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CN, CY, DE, DK,

ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). GDBEN: PIXXD2. APPLICATION: WO 2000-0323975 20000901. PRIORITY: US 1999-0152197 19990901; US 1999-0153251 19991007.

AB The present invention comprises a method of purifying **platelet** membrane **glycoprotein VI (GPVI)**, **GPVI** peptides, cDNA and protein sequence, and methods for using **GPVI** and **antibodies** directed against **GPVI**. It was shown that the extracellular domain of **GPVI** has potent anti-thrombotic activity. The invention comprises methods of inhibiting thrombosis by inhibiting **platelet** aggregation or **platelet** activation using **antibodies** directed against **GPVI**, or **GPVI** protein, in particular, the extracellular domain of **GPVI**.

L6 ANSWER 4 OF 21 EMBLUS COPYRIGHT 2002 AAS

2001:12818 Document No. 134:3175 **Glycoprotein VI** cDNA and protein from human and murine blood **platelets** and their diagnostic and therapeutic applications. Bushfield, Samantha T.; Villelail, Jean-Luc; Janicot-Bennis, Martine; Vainchenet, William; Gill, Davinder Singh; Qian, Ming; Dorian, Kenneth; Hillman, Millennium Pharmaceuticals, Inc., USA). PCT Int. Appl. WO 2001/03141 A1 2001104, 27 pp. DESIGNATED STATES: W: AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DO, EE, ES, FI, GB, GR, HU, ID, IL, IN, IS, JP, KR, KG, KP, KZ, LC, LR, LS, LT, LU, LV, MA, MD, MG, MP, MN, MJ, ME, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, SL, TJ, TM, TR, TT, UA, US, VE, VN, YU, ZA, ZW, FI, AB, BY, EG, EC, MD, RU, TD, TN; RW: AT, BE, BF, BI, CF, CG, CH, CI, IL, IT, DE, DK, ES, FI, FR, GA, GF, GR, IE, IT, LU, IL, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). GDBEN: PIXXD1. APPLICATION: WO 2000-0323975 20000901. PRIORITY: US 1999-0154463 19990630; US 1999-0154324 19991106; US 2000-033387 20000214.

AB The invention provides isolated cDNA mol. and polypeptide mol. that encode human and murine **glycoprotein VI**, a **platelet** membrane glycoprotein that is involved **platelet** -collagen interactions. The protein initially designated TANGO 268 represents the **platelet**-expressed collagen receptor **glycoprotein VI (GPVI)** based on the following evidence: (1) the glycosylated mol. wt. of TANGO 268 and **GPVI** are identical or similar; (2) both are recognized by anti-**GPVI** **antibodies** and bind to convulxin; (3) both are preferentially expressed in megakaryocytic cells; (4) both is predicted to have a single N-glycosylation site; (5) the mol. mass of **GPVI** upon N- and O-linked glycosylation is approx. 62 kDa, that of **GPVI**; (6) two Ig-like domains in TANGO 268 indicates interaction with FcR.gamma.; (7) the absence of a large intracytoplasmic tail suggests that this membrane-bound glycoprotein has no signaling role but assoc. with another member of the Ig family; and (8) TANGO 268 has a charged arginine residue in the transmembrane domain which is also predicted to be present in **GPVI**. The human gene for **GPVI** was mapped on radiation hybrid panels to the long arm of chromosome 19, in the region 19q13, syntenic to mouse chromosome 7. The invention also provides antisense nucleic acid mol., expression vectors contg. the nucleic acid mol. of the invention, host cells into which the expression vectors have been introduced, and non-human transgenic animals in which a nucleic acid mol. of the invention has been introduced or disrupted. The invention still further provides isolated polypeptides, fusion polypeptides, antigenic peptides and **antibodies**. Diagnostic, screening and therapeutic methods utilizing compns. of the invention are also provided.

L6 ANSWER 5 OF 21

MEDLINE

DUPLICATE 2

2001436546 Document Number: 11559356. Pubmed ID: 11544165. A novel viper venom metalloproteinase, alborhagin, is an agonist at the **platelet** collagen receptor **GPVI**. Andrews R K; Gardiner E E; Asazuma N;

Berlanga O; Tulaine D; Nieswandt B; Smith A I; Berndt M C; Watson S P.
Hazel and Pip Appel Vascular Biology Laboratory and the Peptide Biology
Laboratory, Baker Medical Research Institute, Melbourne 3008, Australia..
rkandrews@hotmail.com . JOURNAL OF BIOLOGICAL CHEMISTRY, (2001 Jul 27)
276 (20) 18992-7. Journal code: 1985111R. ISSN: 0021-9258. Pub. country:
United States. Language: English.

AB The interaction of **platelet** membrane **glycoprotein**

VI (GPVI) with collagen can initiate

platelet physiological thrombus formation. The viper venom C-type lectin
family proteins convulxin and alboraggregin-A activate **platelets**
by interacting with GPVI. In this study, we isolated from
white-lipped tree viper (*Trimeresurus albolabris*) venom, alborhagin, which
is functionally related to convulxin because it activates
platelets but is structurally different and related to venom
metalloproteinases. Alborhagin-induced **platelet** aggregation
EC50, 4.5 ng/ml was inhibitable by an anti alphaIIb beta3
antibody, GR304, and the Src family kinase inhibitor PP1,
suggesting that alborhagin activates **platelets**, leading to
alphaIIb beta3-dependent aggregation. Additional evidence suggested that,
like convulxin, alborhagin activated **platelets** by a mechanism
involving GPVI. First, alborhagin- and convulxin-treated
platelets showed a similar tyrosine phosphorylation pattern,
including a similar level of phospholipase Cgamma1 phosphorylation.
Second, alborhagin induced GPVI-dependent responses in
GPVI-transfected K562 and Jurkat cells. Third,
alborhagin-dependent aggregation of mouse **platelets** was
inhibited by the anti-GPVI monoclonal antibody
JAQ1. Alborhagin had minimal effect on convulxin binding to GPVI
-expressing cells, indicating that these venom proteins may recognize
distinct binding sites. Characterization of alborhagin as a GPVI
agonist that is structurally distinct from convulxin demonstrates the
versatility of snake venom toxins and provides a novel probe for
GPVI-dependent **platelet** activation.

L6 ANSWER 6 OF 11 MEDLINE DUPLICATE 4
2001393776 Document Number: 21529121. PubMed ID: 11351023. Rhodocytin
aggregates **platelets** lacking alpha 2 beta 1 integrin,
glycoprotein VI, and the ligand-binding domain of
glycoprotein Ib/alpha. Bergmeier W; Bouvard D; Ehle T A; Mohtari-Nejad R;
Schulte W; Timpone J H; Brackmann C; Essner R; Nieswandt B. (Department
of Molecular Oncology, General Surgery, Witten/Herdecke University,
Artenbergstr. 13, Haus 10, 42117 Wuppertal, Germany. JOURNAL OF
BIOLOGICAL CHEMISTRY, (2001 Jul 6) 276 (13) 25121-6. Journal code:
1985111R. ISSN: 0021-9258. Pub. country: United States. Language: English.

AB Although alpha(2) beta(1) integrin (glycoprotein Ia/IIb) has been
established as a **platelet** collagen receptor, its role in
collagen-induced **platelet** activation has been controversial.
Recently, it has been demonstrated that rhodocytin (also termed aggrexin),
a snake venom toxin purified from the venom of *Calloselasma rhodostoma*,
induces **platelet** activation that can be blocked by
monoclonal antibodies against alpha(2) beta(1) integrin.
This finding suggested that clustering of alpha(2) beta(1) integrin by
rhodocytin is sufficient to induce **platelet** activation and led
to the hypothesis that collagen may activate **platelets** by a
similar mechanism. In contrast to these findings, we provided evidence
that rhodocytin does not bind to alpha(2) beta(1) integrin. Here we show
that the Cre/LoxP-mediated loss of beta(1) integrin on mouse
platelets has no effect on rhodocytin-induced **platelet**
activation, excluding an essential role of alpha(2) beta(1) integrin in
this process. Furthermore, proteolytic cleavage of the 45-kDa N-terminal
domain of glycoprotein (GP) Ib/alpha either on normal or on beta(1)-null
platelets had no significant effect on rhodocytin-induced
platelet activation. Moreover, mouse **platelets** lacking

both alpha(2)beta(1) integrin and the activating collagen receptor **GPVI** responded normally to rhodocytin. Finally, even after additional proteolytic removal of the 45-kDa N-terminal domain of GPIIb/alpha rhodocytin induced aggregation of these **platelets**. These results demonstrate that rhodocytin induces **platelet** activation by mechanisms that are fundamentally different from those induced by collagen.

L6 ANSWER 7 OF 21 MEDLINE DUPLICATE 5
200130478 Document Number: 21293034. PubMed ID: 11237424. Aggretin, a heterodimeric C type lectin from Calloselasma rhodostoma (malayan pit viper), stimulates **platelets** by binding to alpha 2beta 1 integrin and glycoprotein Ib, activating Src and phospholipase Cgamma 2, but does not involve the **glycoprotein VI/IIb** receptor gamma chain collagen receptor. Navdaev A; Clementson J M; Polgar J; Kehrel B E; Glauber M; Haeguenat E; Wells T N; Clementson H J. (Theodor Kocher Institute, University of Berne, Freiestrasse 1, CH-3012 Berne, Switzerland.) JOURNAL OF BIOLOGICAL CHEMISTRY, (2001 Jun 15) 276 (24) 2681-9. Journal code: 2985111R. ISSN: 0021-9158. Pub. country: United States. Language: English.

AB Aggretin, a potent **platelet** activator, was isolated from Calloselasma rhodostoma venom, and 30-amino acid N-terminal sequences of both subunits were determined. Aggretin belongs to the heterodimeric snake C-type lectin family and is thought to activate **platelets** by binding to **platelet** glycoprotein alpha(2)beta(1). We now show that binding to glycoprotein (GP) Ib is also required. Aggretin-induced **platelet** activation was inhibited by a **monoclonal antibody** to GPIb as well as by **antibodies** to alpha(2)beta(1). Binding of both of these **platelet** receptors to aggretin was confirmed by affinity chromatography. No binding of other major **platelet** membrane glycoproteins, in particular **GPVI**, to aggretin was detected. Aggretin also activates **platelets** from Fc receptor gamma chain Fc gamma C-deficient mice to a greater extent than those from normal control mice, showing that it does not use the **GPVI** Fc gamma pathway. **Platelets** from Fc gamma C-deficient mice expressed fibrinogen receptors normally in response to collagen, although they did not aggregate, indicating that these **platelets** may partly compensate via other receptors including alpha(2)beta(1) or GPIb for the lack of the Fc gamma pathway. Signaling by aggretin involves a dose-dependent lag phase followed by rapid tyrosine phosphorylation of a number of proteins. Among these are p72(SYK), p132-FAK, and PLCgamma2, whereas, in comparison with collagen and convulxin, the Fc gamma subunit neither is phosphorylated nor coprecipitates with p72(SYK). This supports an independent, GPIb- and integrin-based pathway for activation of p72(SYK) not involving the Fc gamma receptor.

L6 ANSWER 8 OF 21 MEDLINE DUPLICATE 6
200137088 Document Number: 21226781. PubMed ID: 11278467. Expression and function of the collagen receptor **GPVI** during megakaryocyte maturation. Lagrèze-Lak-Hal A H; Debile N; Hingbury G; Lecut C; Le Couedic J B; Villeviel J L; Jandrot-Perrus M; Vainchenker W. (INSERM E9907, Faculte Xavier Bichat, 75370 Paris Cedex 13, Paris, France.) JOURNAL OF BIOLOGICAL CHEMISTRY, (2001 May 4) 276 (18) 15316-25. Journal code: 2985111R. ISSN: 0021-9258. Pub. country: United States. Language: English.

AB In this report, the expression and function of the **platelet** collagen receptor **glycoprotein VI (GPVI)** were studied in human megakaryocytes during differentiation and maturation of mobilized blood and cord blood derived CD34(+) cells. By flow cytometry, using an anti-**GPVI monoclonal antibody** or convulxin, a **GPVI**-specific ligand, **GPVI** was detected only on CD41(+) cells including some CD41(+)/CD34(+) cells, suggesting expression at a stage of differentiation

similar to CD41. These results were confirmed at the mRNA level using reverse transcription-polymerase chain reaction. **GPVI** expression was low during megakaryocytic differentiation but increased in the more mature megakaryocytes (CD41(high)). As in **platelets**, megakaryocyte **GPVI** associates with the Fc receptor gamma chain (FcRgamma). The FcR gamma chain was detected at the RNA and protein level at all stages of megakaryocyte maturation preceding the expression of **GPVI**. The other collagen receptor, alpha 2(beta 1) integrin (CD49b/CD13), had a pattern of expression similar to **GPVI**. Megakaryocytic **GPVI** was recognized as a 55-kDa protein by immunoblotting and ligand blotting, and thus it presented a slightly lower apparent molecular mass than **platelet GPVI** (58 kDa). Megakaryocytes began to adhere to immobilized convulxin via **GPVI** after only 3-10 days of culture, at a time when megakaryocytes were maturing. At this stage of maturation, they also adhered to immobilized collagen by alpha 2(beta 1) integrin-dependent and -independent mechanisms. Convulxin induced a very similar pattern of protein tyrosine phosphorylation in megakaryocytes and **platelets** including Src, FcRgamma, and PLC gamma3. Our results showed that **GPVI** is expressed early during megakaryocytic differentiation but functionally allows megakaryocyte adherence to collagen only at late stages of differentiation when its expression increases.

- L6 ANSWER 9 OF 21 MEDLINE DUPLICATE 7
 20013593 Document Number: 2124682 PubMed ID: 11384091 Evidence for cross-talk between **glycoprotein VI** and Gi-coupled receptors during collagen-induced **platelet** aggregation. Nieswandt B; Bergmeier W; Eskly A; Schulte V; Ohlmann P; Cazenave J P; Sirmuhl H; Offermanns S; Sachet C. (Department of Molecular Oncology, General Surgery, Witten-Herdecke University, Arrenbergstrasse 20, 42117 Wuppertal, Germany.. nieswandt@klinikum-wuppertal.de) . BLOOD, (2001 Jun 15) 47: 13: 3129-38. Journal Code: 7601500. ISSN: 0006-4971. Pub. Country: United States. Language: English.
- AB Collagen-induced **platelet** aggregation is a complex process and involves synergistic action of integrins, immunoglobulin (Ig)-like receptors, G-protein-coupled receptors and their ligands, most importantly collagen itself, thromboxane A2 (TXA2), and adenosine diphosphate (ADP). The precise role of each of these receptor systems in the overall processes of activation and aggregation, however, is still poorly defined. Among the collagen receptors expressed on **platelets**, glycoprotein (GP) VI has been identified to play a crucial role in collagen-induced activation. **GPVI** is associated with the FcRgamma chain, which serves as the signal transducing unit of the receptor complex. It is well known that clustering of **GPVI** by highly specific agonists results in **platelet** activation and irreversible aggregation, but it is unclear whether collagen has the same effect on the receptor. This study shows that **platelets** from Galpha3-deficient mice, despite their severely impaired response to collagen, normally aggregate on clustering of **GPVI**, suggesting this not to be the principal mechanism by which collagen activates **platelets**. On the other hand, dimerization of **GPVI** by a monoclonal antibody (JAQ1), which by itself did not induce aggregation, provided a sufficient stimulus to potentiate **platelet** responses to Gi-coupled, but not Gq-coupled, agonists. The combination of JAQ1 and adrenaline or ADP, but not serotonin, resulted in alpha 2(beta 1)-dependent aggregation that occurred without intracellular calcium mobilization and shape change in the absence of Galpha3 or the P2Y1 receptor. Together, these results provide evidence for a cross-talk between (dimerized) **GPVI** and Gi-coupled receptors during collagen-induced **platelet** aggregation. (Blood. 2001;97:3129-3135)

Glycoprotein VI but not **alpha2beta1** integrin is essential for **platelet** interaction with collagen. Nieswandt B; Brakebusch C; Bergmeier W; Schulte M; Boeckler D; Mokhtari-Nejad R; Lindhout T; Heemskerk J W; Siengbühl H; Fässler R. (Department of Molecular Oncology, General Surgery, Witten-Herdecke University, 42117 Wuppertal, Germany.. nieswandt@klinikum-wuppertal.de . EMBO JOURNAL, (2001 May 1) 20 (9) 2123-30. Journal code: EMBOJ054. ISSN: 0950-4230. Pub. country: England; United Kingdom. Language: English.

- AB **Platelet** adhesion and activation by components of the extracellular matrix are crucial to arrest post-traumatic bleeding, but can also harm tissue by occluding diseased vessels. Integrin **alpha2beta1** is thought to be essential for **platelet** adhesion to subendothelial collagen, facilitating subsequent interactions with the activating **platelet** collagen receptor, **glycoprotein VI** (GPVI). Here we show that Grelxin-mediated loss of **beta1** integrin on **platelets** has no significant effect on the bleeding time in mice. Aggregation of **beta1**-null **platelets** to native fibrillar collagen is delayed, but not reduced, whereas aggregation to enzymatically digested soluble collagen is abolished. Furthermore, **beta1**-null **platelets** adhere to fibrillar, but not soluble collagen under static as well as low (10 s⁻¹) and high (1000 s⁻¹) shear flow conditions, probably through binding of **alpha11bbeta3** to von Willebrand factor. On the other hand, we show that **platelets** lacking GPVI can not activate integrins and consequently fail to adhere to and aggregate on fibrillar as well as soluble collagen. These data show that GPVI plays the central role in **platelet** collagen interactions by activating different adhesive receptors, including **alpha2beta1** integrin, which strengthens adhesion without being essential.

L6 ANSWER 11 OF 31. EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.

2001406734 EMBASE Bilinexin, a snake C-type lectin from Agkistrodon bilineatus venom agglutinates **platelets** via GPIb and **alpha. 2 .beta. 1**. Du X.-Y.; Nardiev A.; Clemetson J.M.; Magnenat E.; Wells T.M.T.; Clemetson K.J.. Dr. K.J. Clemetson, Theodor Kocher Institute, University of Berne, Freie Strasse 1, CH-3012 Berne, Switzerland. clemetson@ki.kobe.ch. Thrombosis and Haemostasis 86/5 (1977-1286) 2001. Refs: 25.

ISSN: 0340-6245. CODEN: THRODQ. Pub. Country: Germany. Language: English. Summary Language: English.

- AB A new snake protein, named bilinexin, has been purified from Agkistrodon bilineatus venom by ion-exchange chromatography and gel filtration chromatography. Under non-reducing conditions it has a mass of 110 kDa protein on SDS-PAGE. On reduction, it can be separated into five subunits with masses in the range 18-20 kDa. The N-terminal sequences of these subunits are very similar to those of convulxin or the albogaggrins, identifying bilinexin as a new member of the snake C-type lectin family, unusual in having multiple subunits. Bilinexin agglutinates fixed **platelets**, washed **platelets** and **platelet** rich plasma (PRP) without obvious activation (shape change) as confirmed by light microscope examination. Both inhibitory and binding studies indicate that **antibodies** against **alpha. 2 .beta. 1** inhibit not only **platelet** agglutination induced by bilinexin, but also bilinexin binding to **platelets**. WM6d, a monoclonal anti-GPIb.alpha. **antibody**, completely inhibits **platelet** agglutination induced by bilinexin, and polyclonal **antibodies** against GPIb.alpha. prevent its binding to **platelets**. However, neither convulxin, polyclonal anti-GPVI **antibodies**, nor GPIIb/IIIa inhibitors affect its binding to and agglutination of **platelets**. Bilinexin neither activates GPIIb/IIIa integrin on **platelets** nor induces tyrosine phosphorylation of **platelet**

proteins, nor increases intracellular Ca^{2+} in **platelets**. Like alboaggregin B, bilirubin agglutinates **platelets**, which makes it a good tool to investigate the differences in mechanism between snake C-type lectins causing **platelet** agglutination and those that induce full activation.

L6 ANSWER 12 OF 11 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
2002:161488 Document No.: EREV200200261401. Expression of **GPVI** alone confers collagen signaling in RBL-2H3 cells but inactivation of both **GPVI** and $\alpha 2 \beta 1$ is required to inhibit the collagen response of human **platelets**. Chen, Hong (1); Locke, Darren (1); Liu, Changrong (1); Liu, Ying (1); Fahn, Mark L. (1). *J Mol Cell Cardiol*, (November 16, 2001) Vol. 33, No. 1, Part 1, pp. 786a-787a. <http://www.bloodjournal.org/>. print. Meeting Info.: 43rd Annual Meeting of the American Society of Hematology, Part 1, Orlando, Florida, USA December 7-11, 2001 ISSN: 1555-4071. Language: English.

AB The responses of **platelets** to collagen are primary events in arterial thrombosis and are believed to be mediated by two receptors, **GPVI**-Fc gamma and the integrin $\alpha 2 \beta 1$. To determine the role of human **GPVI** we have expressed **GPVI** in RBL-2H3 cells, a mast cell line which expresses abundant Fc gamma but no known collagen receptors, and developed blocking **monoclonal antibodies** to human **GPVI**. These experiments revealed for the first time that RBL-2H3 cells expressing high levels of **GPVI** are capable of both adhesion and calcium signaling in response to fibrillar collagen. Cells expressing lower levels of **GPVI** exhibited adhesion but not signaling, or failed to respond to collagen. Quantitation of **GPVI** receptor density on the surface of **GPVI**-expressing RBL-2H3 cells using 125I-labelled anti-**GPVI** **monoclonal antibody** revealed that the **GPVI** receptor density on high expressing clones is equivalent to that found in human **platelets** (approximately 140 receptors/**platelet**). To test whether **GPVI** is required for collagen responses in human **platelets** we developed a **monoclonal antibody**, 11A12, which blocks calcium signaling in response to collagen but not the **GPVI** agonist convulxin in RBL-2H3 cells. 50 $\mu\text{g/ml}$ 11A12 had a small inhibitory effect on **platelet** aggregation induced by low (1 $\mu\text{g/ml}$) but not high concentrations of collagen (10 and 50 $\mu\text{g/ml}$). A similar small inhibitory effect was observed with the $\alpha 2 \beta 1$ -blocking **antibody** 6F1 used at the same concentration. Strikingly, a combination of 11A12 and 6F1 virtually ablated **platelet** aggregation in response to collagen (10 and 50 $\mu\text{g/ml}$). Our results suggest that (1) **GPVI** is sufficient for both adhesive and signaling responses to collagen; (2) **GPVI**-mediated collagen responses are receptor-density dependent; (3) inhibition of collagen stimulated aggregation of human **platelets** requires inhibition of both **GPVI** and $\alpha 2 \beta 1$. Experiments are currently underway to determine whether the synergistic effect of blocking both $\alpha 2 \beta 1$ and **GPVI** is due to inhibition of $\alpha 2 \beta 1$ -dependent collagen interaction with **GPVI**, **GPVI**-dependent activation of the $\alpha 2 \beta 1$ integrin or to simultaneous inhibition of intracellular signaling by both receptors.

L6 ANSWER 11 OF 21 MEDLINE DUPLICATE 8
2001216216 Document Number: 2116216. PubMed ID: 1133698. Long-term antithrombotic protection by in vivo depletion of **platelet glycoprotein VI** in mice. Nieswandt B; Schulte M; Bergmeier W; Mokhtari-Nejad R; Ruckebardt K; Tazewski J P; Ohlmann P; Gachet C; Zirngibl H. (Department of Molecular Oncology, General Surgery, Witten/Herdecke University, 42117 Wuppertal, Germany. nieswandt@klinikum-wuppertal.de). *JOURNAL OF EXPERIMENTAL MEDICINE*, (2001 Feb 19) 193 (4) 459-69. Journal code: 2965109X. ISSN: 0022-1007. Pub.

country: United States. Language: English.

AB Coronary artery thrombosis is often initiated by abrupt disruption of the atherosclerotic plaque and activation of **platelets** on the subendothelial layers in the disrupted plaque. The extracellular matrix protein collagen is the most thrombogenic constituent of the subendothelial layer; therefore, a selective inhibition of the collagen activation pathway in **platelets** may provide strong antithrombotic protection while preserving other **platelet** functions. Here we demonstrate that treatment of mice with a **monoclonal antibody** against the activating **platelet** collagen receptor **glycoprotein VI** (**GPVI**; **JAQ1**) results in specific depletion of the receptor from circulating **platelets** and abolished responses of these cells to collagen and collagen-related peptides (CRPs). **JAQ1**-treated mice were completely protected for at least 2 wk against lethal thromboembolism induced by infusion of a mixture of collagen (1.8 mg/kg) and epinephrine (6 microg/ml). The tail bleeding times in **JAQ1**-treated mice were only moderately increased compared with control mice probably because the treatment did not affect **platelet** activation by other agonists such as adenosine diphosphate or phorbol myristate acetate. These results suggest that **GPVI** might become a target for long-term prophylaxis of ischemic cardiovascular diseases and provide the first evidence that it is possible to specifically deplete an activating glycoprotein receptor from circulating **platelets** *in vivo*.

L6 ANSWER 14 OF 21 MEDLINE DUPLICATE
200111065 Document Number: 10576376. PubMed ID: 11036073. Evidence for two distinct epitopes within collagen I for activation of murine **platelets**. Schulte V; Schnell D; Bergmeier W; Klingl H; Watson S E; Nieswandt B. Department of Molecular Oncology, General Surgery, Witten-Herdecke University, 42119 Wuppertal, Germany. JOURNAL OF BIOLOGICAL CHEMISTRY, (2001 Jan 5) 276 (1) 364-8. Journal code: 2985121R. ISSN: 0021-9258. Pub. country: United States. Language: English.

AB It has recently been shown that the **monoclonal antibody** **JAQ1** to murine **glycoprotein VI** (**GPVI**) can cause aggregation of mouse **platelets** upon **antibody** cross-linking and that collagen-induced **platelet** aggregation can be inhibited by preincubation of **platelets** with **JAQ1** in the absence of cross-linking. Nieswandt, B., Bergmeier, W., Schulte, V., Ruckebardt, E., Jesner, J. E., and Klingl, H. (2000) J. Biol. Chem. 275, 21991-24001. In the present study, we have shown that cross-linking of **GPVI** by **JAQ1** results in tyrosine phosphorylation of the same profile of proteins as that induced by collagen, including the Fc receptor (FcR) gamma-chain, Src, LAT, SLP-76, and phospholipase C gamma 2. In contrast, **platelet** aggregation and tyrosine phosphorylation of these proteins were inhibited when murine **platelets** were preincubated with **JAQ1** in the absence of cross-linking and were subsequently stimulated with a collagen-related peptide (CRP) that is specific for **GPVI** and low concentrations of collagen. However, at higher concentrations of collagen, but not CRP, aggregation of **platelets** and tyrosine phosphorylation of the above proteins (except for the adapter LAT) is re-established despite the presence of **JAQ1**. These observations suggest that a second activatory binding site, which is distinct from the CRP binding site on **GPVI** on mouse **platelets**, is occupied in the presence of high concentrations of collagen. Although this could be a second site on **GPVI** that is activated by a novel motif within the collagen molecule, the absence of LAT phosphorylation in response to collagen in the presence of **JAQ1** suggests that this is more likely to be caused by activation of a second receptor that is also coupled to the FcR gamma-chain. The possibility that this response is mediated by a receptor that is not coupled to FcR gamma-chain is excluded on the grounds that aggregation is absent in **platelets** from FcR gamma-chain-deficient mice.

receptor **glycoprotein VI** **GPVI** signals

through lipid rafts in a Fc gamma-dependent manner. Locke, Darren (1); Chen, Hong (1); Liu, Chang-Dong (1); Kahn, Mark L. (1). (1) Molecular Cardiology, University of Pennsylvania, Philadelphia, PA USA. Blood, (November 16, 2001) Vol. 98, No. 21 Part 1, pp. 25a.

<http://www.bloodjournal.org/>. print. Meeting Info.: 43rd Annual Meeting of the American Society of Hematology, Part 1 Orlando, Florida, USA December 01-01, 2001 ISSN: 0006-4971. Language: English.

AB The **platelet** collagen receptor **GPVI** signals through the immunoreceptor tyrosine activation motif (ITAM) of its co-receptor Fc gamma using many of the same downstream signaling proteins as T cell, B cell and Fc receptors. Signaling by these immune receptors is believed to proceed from receptor clustering to ITAM tyrosine phosphorylation by the src family kinases Fyn and Lyn and subsequent activation of the tyrosine kinases Syk or JAK-70. Activation of immune receptors results in receptor movement to cholesterol-rich areas of the cell membrane known as lipid rafts that are enriched in Fyn, Lyn and the transmembrane adaptor protein IAT and are defined by their resistance to solubilization by non-ionic detergents. To determine whether activation of **GPVI** results in receptor movement to lipid rafts we expressed **GPVI** in RBL-2H3 cells, a mast cell line which expresses abundant Fc gamma but no known collagen receptors. Activation of **GPVI** with the agonist concanavalin resulted in a rapid, transient movement of **GPVI** receptors to lipid rafts, a response which was also seen with activation of endogenous Fc epsilon receptors which also couple to Fc gamma. The mechanism by which immune receptor activation results in receptor movement to lipid rafts is unknown. To determine the contribution of Fc gamma for **GPVI** movement to lipid rafts we examined the behavior of **GPVI** R27L, a previously characterized mutant **GPVI** receptor in which a single amino acid substitution results in loss of Fc gamma coupling and intracellular signaling despite normal surface expression. **GPVI** R27L binds CVX but does not move to lipid rafts following ligand binding, suggesting that **GPVI** receptor movement to lipid rafts is mediated by the Fc gamma chain. The role of lipid rafts in **platelet** signaling by **GPVI** and other receptors has not been defined. Using a novel anti-**GPVI** monoclonal antibody, HY101, we have isolated lipid rafts from human **platelets** and shown that, like **GPVI** -expressing RBL-2H3 cells, **platelet** stimulation of **GPVI** by concanavalin results in the transient movement of **GPVI** to lipid rafts. Our results demonstrate that 1) during **GPVI** signaling the receptor moves to lipid rafts in both RBL-2H3 cells and in human **platelets**, and 2) **GPVI** movement to lipid rafts following ligand binding is driven by associated Fc gamma chain and is not a simple consequence of ligand-induced receptor clustering. Studies are presently underway to determine whether **GPVI**-Fc gamma movement to lipid rafts is required for ITAM phosphorylation or vice-versa and to better define the role of lipid rafts for signaling by collagen in human **platelets**.

function of the mouse collagen receptor **glycoprotein VI**

is strictly dependent on its association with the Fc gamma chain.

Nieswandt B; Bergmeier W; Schulte V; Ruckebusch K; Gessner J E; Zirngibl H. (Department of Molecular Oncology, General Surgery, University of Witten-Herdecke, 42233 Wuppertal, Germany.. nieswandt@klinikum-wuppertal.de)

JOURNAL OF BIOLOGICAL CHEMISTRY, (2000 Aug 4) 275 (11) 23998-4002.

Journal code: 2085121R. ISSN: 0021-9258. Pub. country: United States.

Language: English.

AB **Platelet glycoprotein (GP) VI** has been proposed as the major collagen receptor for activation of human **platelets**. Human **GPVI** belongs to the immunoglobulin superfamily and is noncovalently associated with the α IIb β 3 chain that is involved in signaling through the receptor. In mice, similar mechanisms seem to exist as **platelets** from α IIb β 3 chain-deficient mice do not aggregate in response to collagen. However, the activating collagen receptor on mouse **platelets** has not been definitively identified. In the current study we examined the function and in vivo expression of **GPVI** in control and α IIb β 3 chain-deficient mice with the first **monoclonal antibody** against **GPVI** (JAQ1). On wild type **platelets**, JAQ1 inhibited **platelet** aggregation induced by collagen but not PMA or thrombin. Cross-linking of bound JAQ1, on the other hand, induced aggregation of wild type but not α IIb β 3 chain-deficient **platelets**. JAQ1 stained **platelets** and megakaryocytes from wild type but not α IIb β 3 chain-deficient mice. Furthermore, JAQ1 recognized **GPVI** (approximately 6-8 kDa) in immunoprecipitation and Western blot experiments with wild type but not α IIb β 3 chain-deficient **platelets**. These results strongly suggest that **GPVI** is the collagen receptor responsible for **platelet** activation in mice and demonstrate that the association with the α IIb β 3 chain is critical for its expression and function.

L6 ANSWER 11 OF 21 BIOSIS COPYRIGHT © 2002 BIOLOGICAL ABSTRACTS INC.
 2001:41199 Document No.: PRE20011091588. Long-term antithrombotic protection by irreversible inactivation of **platelet glycoprotein VI** in mice. Nieswandt, Bernhard (1); Schulze, Valerie (1); Bernmeier, Wolfgang (1); Mochlyar-Negad, Rahee (1); Takemura, Jean R.; Radner, Christian; Zirngibl, Hubert (1). (1) Molecular Oncology, Witten Herderschee University, Wuppertal Germany. Blood, (November 16, 2000) Vol. 86, No. 11 Part 1, pp. 2694. print. Meeting Info.: 42nd Annual Meeting of the American Society of Hematology San Francisco, California, USA December 1-05, 2000 American Society of Hematology. ISSN: 0006-4713. Language: English. Summary Language: English.

AB Coronary artery thrombosis is often initiated by abrupt disruption of the atherosclerotic plaque followed by deposition and activation of **platelets** on the subendothelial layers in the disrupted plaque. Because the extracellular matrix protein collagen is the most thrombogenic constituent of the subendothelial layer, a selective inhibition of the collagen activation pathway in **platelets** may provide strong antithrombotic protection while preserving other **platelet** functions. Growing evidence suggests that **platelet** glycoprotein GPVI is the major collagen receptor for **platelet** activation making this receptor a good candidate for such a specific inhibition. In the current study, we have investigated the antithrombotic effects of the first **monoclonal antibody** mAb against mouse **GPVI** (JAQ1, Nieswandt et al; 2001, J Biol Chem, 276(31):23498-23502). Injection of 100 μ g JAQ1 only had mild and transient effects on **platelet** counts with a maximum drop of approximately $34 \pm 7.1\%$ on day 1 and a return to normal after 2-3 days. JAQ1 pretreated mice were completely protected against lethal thromboembolism induced by infusion of a mixture of collagen (5.8 mg/kg) and epinephrine (60 μ g/kg) for at least two weeks: 100% survivors on days 1, 7, and 14 after mAb injection, n=3 per group, 5% survivors in the control group, n=20). Aggregometric and flow cytometric studies demonstrated that **platelets** from JAQ1 treated mice were completely resistant against activation with high concentrations of collagen (up to 100 μ g/ml) and collagen related peptides (up to 100 μ g/ml) ex vivo on days 1, 7, and 14. In JAQ1 treated mice, **GPVI** was not detectable in a Western blot analysis of **platelet** lysates for minimally two weeks, suggesting irreversible inactivation (or degradation) of the receptor on circulating **platelets**. In

contrast to collagen, other agonists, such as ADP or **platelet** aggregating agents, such as PMA induced normal activation and aggregation of these **platelets**. Consequently, the tail bleeding times were only moderately increased in anti-GPVI treated mice compared to control mice on day 2, 7, and 14. These results establish **GPVI** as an attractive target for long-term antithrombotic therapy.

L6 ANSWER 18 OF 11 MEDLINE DUPLICATE 11
19991-7242 Document Number: 99167-43. PubMed ID: 1066433. Signal transduction pathways mediated by glycoprotein IIb/IIIa in human **platelets**: comparison with those of **glycoprotein** VI. Inoue K; Ozaki Y; Satoh K; Wu Y; Yotomi Y; Sano Y; Morita T. Department of Clinical and Laboratory Medicine, Yamaguchi Medical University, Uramoto 1-1-1 Yamaguchi, Yamaguchi, 750-8511, Japan.) BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (1999 Mar 5) 256 (1) 114-21. Journal title: 0882251. ISSN: 0006-294X. PDB. Country: United States. Language: English.

AB Human **platelets** were activated either by glycoprotein (GP) Ia/IIa agonist (rhodocytin) or by a **GPVI** agonist (collagen-related peptide, CRP), and the intracellular signal transduction pathways were compared in the presence of various inhibitors. Rhodocytin isolated from *Callisaurus draconoides* venom was verified as a GPIa/IIa agonist, based on the inhibitory effects of three mAbs directed against GPIa. **Platelet** activation mediated by GPIa/IIa led to overt **platelet** aggregation, elevation of intracellular Ca^{2+} , and tyrosine phosphorylation of several proteins, similar to that of **GPVI**. p72 syk and phospholipase Cgamma1 (PLCgamma1) tyrosine phosphorylation were also observed with GPIa/IIa-mediated **platelet** aggregation, although they peaked slightly later than that of **GPVI**. In contrast to **GPVI**-mediated **platelet** activation, most of these phenomena induced by GPIa/IIa activation were markedly suppressed by acetylsalicylic acid (ASA) or cytochalasin D. These findings suggest that the requirements for thromboxane A2 (TXA2) production and actin polymerization, which are the characteristics of collagen-induced **platelet** activation, are derived from the GPIa/IIa-mediated signal transduction, but not from that of **GPVI**. Copyright 1999 Academic Press.

L6 ANSWER 19 OF 11 RESEARCH COPYRIGHT 2002 ISI (R)
1998: 66583 The Genuine Article R Number: 2M110. Simple collagen-like peptides support **platelet** adhesion under static but not under flow conditions: Interaction via alpha beta 1 and von Willebrand factor with specific sequences in native collagen is a requirement to resist shear forces. Verkleij M W (Reprint); Morton L F; Knight C G; deGroot P G; Barnes M J; Fimia J J. UNIV WIRECHT HOSP, DEPT HAEMATOLOGY, POSTGRADUAL FOR BIOMEMBRANES, P.O. BOX 350, NL-3500 GA UTRECHT, NETHERLANDS (Reprint); STRANDEWAY RES LAB, CAMBRIDGE CB1 4RN, ENGLAND. BLOOD 15 MAY 1998 Vol. 91, No. 10, pp. 3315-3316. Publisher: W B SAUNDERS CO, INDEPENDENCE SQUARE WEST, SUITE CENTER, STE 100, PHILADELPHIA, PA 19106-3698. ISSN: 0006-4971. PDB. Country: NETHERLANDS; ENGLAND. Language: English.
ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS

AB The aim of this study was to define the need for specific collagen sequences and the role of their conformation in **platelet** adhesion to collagen under both static and flow conditions. We recently reported that simple triple-helical collagen-related peptides (CRPs), CRP*(GEP*) (10)GCH*E and CRP*(GPP*) (10)GPP*G (single-letter amino acid code, P = proline; H = hydroxyproline; Morton et al, Biochem J 306:337, 1995) were potent stimulators of **platelet** activation and were able to support the adhesion of gel-filtered **platelets** examined under static condition. The present study investigated whether these same peptides were able to support **platelet** adhesion under more physiologic conditions by examining static adhesion with **platelet**-rich plasma (PRP) and adhesion underflow conditions. In the static

adhesion assay, we observed 20% surface coverage with **platelet** aggregates. In marked contrast, there was a total lack of adhesion under flow conditions examined at shear rates of 50 and 200 s⁻¹). Thus, the interaction of **platelets** with the CRPs is a low-affinity interaction unable on its own to withstand shear forces. However, the addition of CRPs to whole blood, in the presence of 200 mu mol/LD-arginyl-glycyl-L-aspartyl-L-tryptophan (GRGSP) to prevent **platelet** aggregation, caused an inhibition of about 50% of **platelet** adhesion to collagens I and III under flow. These results suggest that the collagen triple helix per se, as defined by these simple collagen sequences, plays an important contributory role in the overall process of adhesion to collagen under flow. The **monoclonal antibody** (1F6D), directed against the alpha 1 subunit of the integrin alpha 2 beta 1, was found to inhibit static **platelet** adhesion to monomeric but not fibrillar collagens I and III. However, under flow conditions, anti-alpha 2 MoAbs (17G07 and 6F1) inhibited adhesion to both monomeric and fibrillar collagens, indicating that alpha 2 beta 1 is essential for adhesion to collagen under flow, independent of collagen conformation, whether monomeric or polymeric. To obtain further insights into the nature of the different adhesive properties of CRPs and native collagen, we investigated the relative importance of von Willebrand factor (vWF) and the integrin alpha 2 beta 1 in **platelet** adhesion to collagen types I and III, using the same shear rate (200 s⁻¹) as used when testing CRPs under flow conditions. Our results, together with recent data of others, support a two step mechanism of **platelet** interaction with collagen under flow conditions. The first step involves adhesion via both the indirect interaction of **platelet** glycoprotein (GP) Ib with collagen mediated by vWF binding to specific vWF recognition sites in collagen and the direct interaction between **platelet** alpha 2 beta 1 and specific alpha 1 beta 1-recognition sites in collagen. This suffices to hold **platelets** at the collagen surface. The second step occurs via another collagen receptor (thought to be GPVI) that binds to simple collagen sequences, required essentially to delineate the collagen triple helix. Recognition of the triple helix leads to strengthening of attachment and **platelet** activation. © 1993 by The American Society of Hematology.

L6 ANSWER 20 OF 11 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 12 1998211166 EMBASE Convolxin-induced **platelet** adhesion and aggregation: Involvement of **glycoproteins VI** and **alpha 2 beta 1**. Jancinet-Perrus H.; Lagnier A.H.; Lelias M.; Ohama M.; Ben G. Dr. M. Jancinet-Perrus, Lab. Recherche Hemostase Thrombose, Faculte de Medecine Xavier Bichat, 75870 Paris Cedex 13, France. Platelets 4(3-4 207-211) 1993.

Refs: 22.

ISSN: 0950-7104. CDDEN: HEMTEEF. Pub. Country: United Kingdom. Language: English. Summary Language: English.

AB The interaction of convolxin (Cvx), a 72-kDa glycoprotein isolated from the venom of *Crotalus durissus terrificus* with human **platelets** has been studied. Cvx at low concentrations (below 100 pM) induced **platelet** aggregation, dense body secretion and intracellular calcium mobilization which indicates that Cvx is a potent activator of human **platelets**. Cvx-induced **platelet** aggregation and secretion was inhibited by 6F1 an anti-integrin .alpha.2.beta.1. **monoclonal antibody** that was without effect on calcium mobilization. Anti-GPVI Fab fragments inhibited aggregation, secretion and calcium mobilization triggered by Cvx. In addition, immobilized Cvx was found to induce divalent cation-independent **platelet** adhesion in a static system. **Platelet** adhesion to Cvx was inhibited by anti-GPVI Fab fragment but not by anti-integrin .alpha.2.beta.1. Cvx was shown to bind to a 57,000 Dalton protein that was identified as GPVI. Altogether, these results

indicate that **GPVI** behaves as a receptor for Cvx, while integrin $\alpha 2 \beta 1$ could play a regulatory role in Cvx-induced **platelet** aggregation. Cvx and collagen interaction with **platelets**, thus appears to share some characteristics but to also have specific properties.

L6 ANSWER 21 OF 21 MEDLINE DUPLICATE 13
 1998001677 Document Number: 98001677. PubMed ID: 9841142. Adhesion and activation of human **platelets** induced by convulxin involve **glycoprotein VI** and integrin $\alpha 2 \beta 1$. Tandrot-Perrus M; Lagrèze A H; Okuma M; Bon C. (Laboratoire de Recherche sur l'Hémostase et la Thrombose, Faculté de Médecine Xavier Bichat, BP 416, 75870 Paris Cedex 18, France.) JOURNAL OF BIOLOGICAL CHEMISTRY, 1997 Oct 24) 272 (48) 27335-41. Journal code: 298121R. ISSN: 0021-9.56. Pub. country: United States. Language: English.

AB We analyzed the interaction of convulxin (Cvx), a 72 kDa protein isolated from the venom of *Crotalus durissus terrificus*, with human **platelets**. Cvx is a potent **platelet** agonist that induces an increase in the intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$), granule exocytosis and aggregation. 125I-labeled Cvx binds specifically and rapidly to **platelets** at binding sites of high and moderate affinity. **Platelets** adhere to immobilized Cvx in a time-dependent but cation-independent manner. **Platelet** exocytosis and aggregation induced by Cvx were inhibited by an anti-integrin $\alpha 2 \beta 1$ monoclonal antibody (6F1) and by the Fab fragments of a polyclonal anti-**glycoprotein VI (GPVI) antibody**. Both the adhesion of **platelets** to Cvx and the Cvx-induced increase in $[Ca^{2+}]_i$ were inhibited by anti-GPVI Fab fragments but not by 6F1. Ligand blotting assay showed that 125I-Cvx binds to a 57-kDa **platelet** protein with an electrophoretic mobility identical to that of GPVI. In addition, we observed the following: (i) 125I-Cvx binds to GPVI immunoprecipitated by the anti-GPVI **antibody** from a **platelet** lysate, and (ii) Cvx inhibits the binding of anti-GPVI IgG to GPVI. Taken together, these results demonstrate that GPVI behaves as a Cvx receptor and that the $\alpha 2 \beta 1$ integrin appears to be involved in the later stages of Cvx-induced **platelet** activation, i.e. exocytosis and aggregation.

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	36.18	36.49
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-1.24	-1.24

STN INTERNATIONAL LOGOFF AT 08:38:32 ON 11 DEC 2002